CORRECTION Open Access

Correction: Risk prediction of 30-day mortality after stroke using machine learning: a nationwide registry-based cohort study

Wenjuan Wang^{1*}, Anthony G. Rudd¹, Yanzhong Wang^{1,2,3}, Vasa Curcin^{1,2,3}, Charles D. Wolfe^{1,2,3}, Niels Peek^{4,5†} and Benjamin Bray^{1†}

Correction: BMC Neurol 22, 195 (2022) https://doi.org/10.1186/s12883-022-02722-1

Following publication of the original article [1], the authors reported an error in Fig. 1. The same error was made in Figure A, C and D in supplemental material. The context was not affected as the error was made when replotting during the revision. We apologise for the error.

The correct Fig. 1 and supplemental material are provided below.

The original article [1] has been updated.

Supplementary Information

The online version contains supplementary material available at https://doi.org/10.1186/s12883-022-02840-w.

Additional file 1.

Author details

¹ School of Population Health & Environmental Sciences, Faculty of Life Science and Medicine, King's College London, London, UK. ²NIHR Biomedical Research Centre, Guy's and St Thomas' NHS Foundation Trust and King's College London, London, UK. ³NIHR Applied Research Collaboration (ARC) South London, London, UK. ⁴Division of Informatics, Imaging and Data Science, School of Health

The original article can be found online at https://doi.org/10.1186/s12883-022-02722-1

[†]Niels Peek and Benjamin Bray are Joint last author

*Correspondence: wenjuan.wang@kcl.ac.uk

¹ School of Population Health & Environmental Sciences, Faculty of Life Science and Medicine, King's College London, London, UK Full list of author information is available at the end of the article

Sciences, University of Manchester, Manchester, UK. ⁵NIHR Manchester Biomedical Research Centre, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK.

Published online: 25 August 2022

Reference

 Wang W, Rudd AG, Wang Y, et al. Risk prediction of 30-day mortality after stroke using machine learning: a nationwide registry-based cohort study. BMC Neurol. 2022;22:195. https://doi.org/10.1186/s12883-022-02722-1.



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Wang et al. BMC Neurology (2022) 22:319 Page 2 of 2

