

ERRATUM

Open Access



Erratum to: A randomized double-blind multi-center trial of hydrogen water for Parkinson's disease: protocol and baseline characteristics

Asako Yoritaka^{1,2*}, Takashi Abe³, Chigumi Ohtsuka⁴, Tetsuya Maeda⁵, Masaaki Hirayama⁶, Hirohisa Watanabe⁷, Hidemoto Saiki⁸, Genko Oyama², Jiro Fukae⁹, Yasushi Shimo², Taku Hatano², Sumihiro Kawajiri¹⁰, Yasuyuki Okuma¹⁰, Yutaka Machida¹¹, Hideto Miwa¹¹, Chikako Suzuki¹², Asuka Kazama¹³, Masahiko Tomiyama¹⁴, Takeshi Kihara¹⁵, Motoyuki Hirasawa¹⁶, Hideki Shimura¹⁷ and Nobutaka Hattori²

Erratum

After publication of the original article [1], it came to the authors' attention that there was an error in the number of male and female participants. The correct number is 90 male and 88 female, not 89 male and 89 female as originally reported.

This error was present in the Methods section of the abstract and in Table 1.

Author details

¹Department of Neurology, Juntendo University Koshigaya Hospital, Fukuroyama 560, Koshigayashi, Saitama 343-0032, Japan. ²Department of Neurology, Juntendo University School of Medicine, Tokyo, Japan. ³Department of Neurology, Abe Neurological Clinic, Iwate, Japan. ⁴Department of Neurology and Gerontology, Iwate Medical University, Iwate, Japan. ⁵Department of Neurology, Research Institute for Brain and Blood Vessels-Akita Hospital, Akita, Japan. ⁶Department of Pathophysiological Laboratory Sciences, Nagoya University Graduate School of Medicine, Aichi, Japan. ⁷Brain and Mind Research Center, Nagoya University Graduate School of Medicine, Aichi, Japan. ⁸Department of Neurology, Kitano Hospital, The Tazuke Kofukai Medical Research Institute, Osaka, Japan. ⁹Department of Neurology, Fukuoka University, Fukuoka, Japan. ¹⁰Department of Neurology, Juntendo University Shizuoka Hospital, Shizuoka, Japan. ¹¹Department of Neurology, Juntendo University Nerima Hospital, Tokyo, Japan. ¹²Department of Diagnostic Radiology, Department of Molecular Medicine and Surgery, Karolinska University Hospital, Karolinska Institute, Stockholm, Sweden. ¹³Nozomi Hospital, Saitama, Japan. ¹⁴Department of Neurology, Aomori Prefectural Central Hospital, Aomori, Japan. ¹⁵Department of Neurology, Rakuwakai Otowa Rehabilitation Hospital, Kyoto, Japan. ¹⁶Department of Neurology, Tokyo Rinkai Hospital, Tokyo, Japan. ¹⁷Department of Neurology, Juntendo University Urayasu Hospital, Chiba, Japan.

Published online: 20 February 2017

Reference

1. Yoritaka A, Abe T, Ohatsuka C, et al. A randomized double-blind multi-center trial of hydrogen water for Parkinson's disease: protocol and baseline characteristics. *BMC Neurol.* 2016;16:66.

* Correspondence: ayori@juntendo.ac.jp

¹Department of Neurology, Juntendo University Koshigaya Hospital, Fukuroyama 560, Koshigayashi, Saitama 343-0032, Japan

²Department of Neurology, Juntendo University School of Medicine, Tokyo, Japan

Table 1 The baseline characteristics of the study participants

	Mean	Std. Deviation
Age	64.2	9.2
Disease duration	6.8	4.5
Sex (male/female) n	90	88
Wearing off (+/-) n	58	120
Dyskinesia (+/-) n	42	136
Levodopa mg	344.1	202.8
Levodopa (+/-) n	154	24
Dopamine agonist levodopa equivalent dose (mg) [12]	166.3	121.1
Total levodopa equivalent dose (mg) [12] ^a	592.0	317.6
Unified Parkinson's Disease Rating Scale		
Part I	0.7	1.2
Part II	5.6	3.9
Part III	15.7	8.4
Part IV	1.7	2.2
Total	23.7	11.8
Modified Hoehn and Yahr stage	2.1	0.6
Parkinson's disease Questionnaire-39		
Mobility	10.4	9.1
Activities of daily living	5.2	5.1
Emotional well-being	6.0	4.4
Stigma	3.2	2.8
Social support	1.4	1.9
Cognitions	4.0	3.0
Communication	1.7	2.2
Body discomfort	2.5	2.6
Total	34.4	24.2

^anot including istradefylline and zonisamide